

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	TSPAN33
<b>Synonyms</b>	PEN; PEN.; TSPAN-33
<b>Description</b>	Human TSPAN33 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q86UF1
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length TSPAN33 protein has a MW of 31.5 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Plays an important role in normal erythropoiesis (By similarity). It has a role in the differentiation of erythroid progenitors (By similarity). Regulates maturation and trafficking of the transmembrane metalloprotease ADAM10 (PubMed:26686862). Negatively regulates ligand-induced Notch activity probably by regulating ADAM10 activity (PubMed:26686862).
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate TSPAN33-Nanodisc 0.2 $\mu$ g Human TSPAN33-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag TSPAN33-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TSPAN33-Nanodisc is 10.31ng/ml.




Figure2. Human TSPAN33-Nanodisc, Flag Tag on SDS-PAGE

