

## PRODUCT INFORMATION

<b>Target</b>	VSIG4
<b>Synonyms</b>	CRlg;Z39IG
<b>Description</b>	Recombinant Human VSIG4 with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9Y279
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc Tag
<b>Molecular Characterization</b>	VSIG4(Arg20-Pro283) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 55.3 kDa after removal of the signal peptide. The apparent molecular mass of VSIG4-hFc is approximately 55-70 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a v-set and immunoglobulin-domain containing protein that is structurally related to the B7 family of immune regulatory proteins. The encoded protein may be a negative regulator of T-cell responses. This protein is also a receptor for the complement component 3 fragments C3b and iC3b. Alternate splicing results in multiple transcript variants. [provided by RefSeq, May 2010]
<b>Usage</b>	Research use only






Figure 1. Human VSIG4 Protein, hFc Tag on SDS-PAGE under reducing condition.

### Human VSIG4,hFc Tagged protein ELISA

0.2 µg of Human VSIG4, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human VSIG4 Protein, hFc Tag(PME100855) can bind Anti-VSIG4 antibody(DMC268), IgG1 Chimeric mAb in a linear range of 3.20-80 ng/mL.

