

**PRODUCT INFORMATION**

<b>Target</b>	ACVR2A
<b>Synonyms</b>	Acvr2; Actr1la; Tactrl1
<b>Description</b>	Recombinant mouse ACVR2A protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P27038
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc tag
<b>Molecular Characterization</b>	Mouse ACVR2A(Ala20-Pro135) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 39.6 kDa after removal of the signal peptide. The apparent molecular mass of mACVR2A-hFc is approximately 35-70 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Enables several functions, including BMP receptor activity; PDZ domain binding activity; and activin binding activity. Acts upstream of or within several processes, including Sertoli cell proliferation; copulation; and embryonic skeletal system development. Located in cell surface. Is expressed in several structures, including central nervous system; early conceptus; gonad; gut; and sensory organ. Used to study Weissenbacher-Zweymuller syndrome. Human ortholog(s) of this gene implicated in colon cancer. Orthologous to human ACVR2A (activin A receptor type 2A). [provided by Alliance of Genome Resources, Apr 2022]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Mouse ACVR2A Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

