

**PRODUCT INFORMATION**

<b>Target</b>	BAFF
<b>Synonyms</b>	B-cell-activating factor;CD257;Tnfsf13b;Baff
<b>Description</b>	Recombinant mouse BAFF protein with N-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9WU72
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Human Fc Tag
<b>Molecular Characterization</b>	hFc(Glu99-Ala330) Mouse BAFF(Ala127-Leu309)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 46.8 kDa after removal of the signal peptide. The apparent molecular mass of hFc-mBAFF is approximately 40-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. TNFSF13/APRIL binds to the same 2 receptors. Together, they form a 2 ligands -2 receptors pathway involved in the stimulation of B- and T-cell function and the regulation of humoral immunity. A third B-cell specific BAFF-receptor (BAFFR/BR3) promotes the survival of mature B-cells and the B-cell response.[UniProtKB/Swiss-Prot Function]
<b>Usage</b>	Research use only






Figure 1. Mouse BAFF Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

