

## PRODUCT INFORMATION

<b>Target</b>	BTLA
<b>Synonyms</b>	BTLA;D272
<b>Description</b>	Recombinant mouse BTLA protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q7TSA3
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	Mouse BTLA(Glu30-Pro176) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 17.4 kDa after removal of the signal peptide. The apparent molecular mass of Mouse-BTLA-His is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Inhibitory receptor on lymphocytes that negatively regulates antigen receptor signaling via PTPN6/SHP-1 and PTPN11/SHP-2 (PubMed:12796776, PubMed:14652006). May interact in cis (on the same cell) or in trans (on other cells) with TNFRSF14 (PubMed:19915044). In cis interactions, appears to play an immune regulatory role inhibiting in trans interactions in naive T cells to maintain a resting state. In trans interactions, can predominate during adaptive immune response to provide survival signals to effector T cells (PubMed:19915044).
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Mouse BTLA Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

