

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Clone ID</b>                         | DM48  |
| <b>Target</b>                           | ACE2  |
| <b>Synonyms</b>                         | ACE-2; ACEH; ACE2   |
| <b>Host Species</b>                     | Rabbit  |
| <b>Description</b>                      | PE-conjugated Anti-ACE2 antibody(DM48); Rabbit mAb  |
| <b>Delivery</b>                         | 3-4 weeks   |
| <b>Uniprot ID</b>                       | Q9BYF1  |
| <b>IgG type</b>                         | Rabbit IgG  |
| <b>Clonality</b>                        | Monoclonal  |
| <b>Reactivity</b>                       | Human   |
| <b>Applications</b>                     | Flow Cyt  |
| <b>Recommended Dilutions</b>            | Flow Cyt 1:100  |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography   |
| <b>Formulation &amp; Reconstitution</b> | N/A   |
| <b>Storage &amp; Shipping</b>           | Store at 2°C-8°C for 6 months   |
| <b>Background</b>                       | The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9; and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function; as well as fertility. In addition; the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63. |
| <b>Usage</b>                            | Research use only   |

