

PRODUCT INFORMATION

Clone ID DM196 CCR8 **Target**

CC-CKR-8; CCR-8; CDw198; CKRL1; CMKBR8; CMKBRL2; CY6; GPRCY6; TER1 **Synonyms**

Host Species

PE-conjugated Anti-CCR8 antibody(DM196); Rabbit mAb **Description**

3-4 weeks

Uniprot ID P51685 IgG type Rabbit IgG Clonality Monoclonal

Reactivity Human **Applications** Flow Cyt

Recommended **Dilutions**

Background

Delivery

Flow Cyt 1:100

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Liquid

☐PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

> This gene encodes a member of the beta chemokine receptor family; which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are important for the migration of various cell types into the inflammatory sites. This receptor protein preferentially expresses in the thymus. I-309; thymus activation-regulated cytokine (TARC) and macrophage inflammatory protein-1 beta (MIP-1 beta) have been identified

as ligands of this receptor. Studies of this receptor and its ligands suggested its role in regulation of monocyte chemotaxis and thymic cell apoptosis. More specifically; this receptor may contribute to the proper positioning of activated T cells within the antigenic challenge sites and specialized areas of lymphoid tissues. This gene is located at the chemokine receptor

> Email: info@dimabio.com Website: www.dimabio.com

gene cluster region.

Usage Research use only

Conjugate PE-conjugated

