

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM191
<b>Target</b>	CD70
<b>Synonyms</b>	CD70;CD27LG;TNFSF7;TNFSF7G;CD27L
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-CD70 antibody(DM191); Rabbit mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	P32970
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27:CD27. It is a surface antigen on activated; but not on resting; T and B lymphocytes. It induces proliferation of costimulated T cells; enhances the generation of cytolytic T cells; and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation; cytotoxic function of natural killer cells; and immunoglobulin synthesis.
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

