

## PRODUCT INFORMATION

<b>Clone ID</b>	DMC390
<b>Target</b>	FOLR2
<b>Synonyms</b>	BETA-HFR; FBP; FBP:PL-1; FOLR1; FR-BETA; FR-P3; FRbeta
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-FOLR2 antibody(DMC390); IgG1 Chimeric mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	P14207
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	<p>The protein encoded by this gene is a member of the folate receptor (FOLR) family; and these genes exist in a cluster on chromosome 11. Members of this gene family have a high affinity for folic acid and for several reduced folic acid derivatives; and they mediate delivery of 5-methyltetrahydrofolate to the interior of cells. This protein has a 68% and 79% sequence homology with the FOLR1 and FOLR3 proteins; respectively. Although this protein was originally thought to be specific to placenta; it can also exist in other tissues; and it may play a role in the transport of methotrexate in synovial macrophages in rheumatoid arthritis patients. Multiple transcript variants that encode the same protein have been found for this gene.</p>
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

