

PRODUCT INFORMATION

Target Farget GPC3GPC3SynonymsDGSX;GTR2-2;MXR7;OCI-5;SDY5;SGB;SGB5;SGB51Host SpeciesRabbitDescriptionChimeric mAbDeliveryUnder DevelopmentUniprot IDP51654IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:00PurificationCurified form cell culture supernatant by affinity chromatographyFormulationStore at 2°C-8°C for 6 monthsStorage & ShippingStore at 2°C-8°C for of monthsStorage & ShippingStore at 2°C-8°C for of mothsBackgroundamembrane-associated protein core substituted with a variable number of hegyptican-feated or yowth regulation. The protein encode by this gene can bind to and inhibit the dipeptidy cytoplasmic membrane via a glycosyl cytoplasmic membrane, also know as Simpson- dysmorpha Syndrome, also know as	Clone ID	163D11
SynonymsDGSX;GTR2-2;MXR7;OCI-5;SDYS;SGB;SGB5;SGB51Host SpeciesRabbitDescriptionPE-conjugated Anti-GPC3 antibody(163D11), IgG1 Chimeric mAbDeliveryUnder DevelopmentUniprot IDP51654IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsCell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan subfate chains, Members of the glypican-related integral membrane typoteoglycan family (GRIPS) cytoplasmic membrane with a glycosyl phosphatidylinositol linkage. These proteins may play arole in the control of cell division and growth regulation. The protein encoded by this gene can bind to and infibit the dipeptidyl peptidase activity of CD26, and it can induce yeptidase activity of CD36, and it can induce yeptidase activity		
Host SpeciesRabbitDescriptionPE-conjugated Anti-GPC3 antibody(163D11), IgG1 Chimeric mAbDeliveryUnder DevelopmentUnjprot IDP51654IgG typeRabbit/Human Ec chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100Purified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionCell surface protein normatographyStorage & ShippingStore at 2°C-8°C for 6 monthsCell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion matagene peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion may peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion may peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion may peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion may flap a nole in the control of cell division and gene can bind to and inhibit the dipeptidyl peptidase activity of CD2, and it can induce appotosis in certain cell types. Deletion mutations in multiple transcript variants. Iprovided by the appetidese activity of CD2, and it can induce appotosis in certain cell types. Deletion mutations in multiple transcript variants. Iprovided by the thepatocellular carcinomal JI, Sci Rep. 2009, Ref	2	
DescriptionPE-conjugated Anti-GPC3 antibody(163D11), IgG1 Chimeric mADDeliveryUnder DevelopmentUniprot IDP51654IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationLiquid]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsCell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chorter of the gene and initigene are associated by this gene can bind to and initibit the dispetidy gene are associated with Simpson-Golabi- Behmel syndrome. Alternative splicing results in multipe transcript variants. Jprovided by RefSeq. Sep 2009] References] Fu Ying, Urban Daniel J, Nail Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma II]. Hepatology, 2019, 70: 563-576. Zhang Yi-Fan, Ho Mitchell-Humanization of high-affinity antibodies targeting diplocan- in the patology. 2019, 70: 563-576. Zhang Yi-Fan, Ho Mitchell-Humanization of high-affinity antibodies targeting diplocan- in thepatology. 2019, 70: 563-576. Zhang Yi-Fan, Ho Mitchell-Humanization of high-affinity antibodies targeting diplocan- in hepatology. 2019, 70: 563-576. Zhang Yi-Fan, Ho Mitchell-Humanization of high-affinity antibodies targeting diplocan- in hepatology. 2019, 70: 563-		
DeliveryUnder DevelopmentUniprot IDP51654IgG typeRabbit/Human Fc chimeric IgG1IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended Dilutions*Flow Cyt 1:100Purified from cell culture supernatant by affinity chromatographyFormulation & recomstitutionLiquid[PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glycican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl peridase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by Refseq. Sep 2009] References] Fu Ying Urban Daniel J.Nain Roger R et al. Glypican-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinomal[]] -Kepatology, 2019, 70: 553-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting dlypican-3 in hepatocellular carcinomal[]] -Kepatology, 2019, 70: 553-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting dlypican-3 in hepatocellular carcinomal[]] -Kepatocellular carcinomal[]] -Kepatology, 2019, 70: 553-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting dlypican-3 in hepatocellular carcinomal[]] -Kepatology, 2	-	PE-conjugated Anti-GPC3 antibody(163D11), IgG1
IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommendedFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsCell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the contol of cell division and growth regulation. The protein encoded by this gene are associated with Simpson-Golabi- Behmel synorme. Alternative splicing results in multiple transcript variants. [provided by Refseq. Sep 2009] ReferenceS[] V Ting. Urban Daniel J, Nani Roger R et al. Glypican-3 Specific Antibody Drug Conjugates Targeting Antibody Drug Conjugates Targeting of Hone Alternative splicing results for Mitchell Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.[]]. Sel Rep. 2016, 6: 3387.8.UsageResearch use onlyConjugatePE-conjugated Alt PIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application to protein agenuencing or reverse engineering attempt is prohibited. We are actively scrutinizing al patent application. The y are all under patent application. Any protein sequencing or reverse engineerin	Delivery	
ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glycican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome, Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009] References[] Lving, Urban Daniel], Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[]] -Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell, Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.[]] -Sci Rep, 2016, 6: 33878.UsageResearch use onlyDIMA DisclaimerPE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to b	Uniprot ID	P51654
ReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glycican-related integral membrane incomed to the cytoplasmic membrane via a glycosyl phosphatigylinosite in core or bubstituted with a variable number of heparan sulfate chains. Members of the glycican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatigylinositel linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome, Alternative splicing results in multiple transcript variants. [provided by MetSeq. Spc 2009] References] Tu Ying, Urban Daniel J. Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[]] -Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.[]] -Sci Rep, 2016, 6: 33878.UsageResearch use onlyDiMA DisclaimerPic-conjugated All DIMA recombinant antibodi	lgG type	Rabbit/Human Fc chimeric IgG1
ApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the diapetidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in multiple transcript variants. [Drovided by RefSeq. Sep 2009] References] Fu Ying.Urban Daniel J.Nani Roger R et al. Glypicangeting respectival carinoma. [J] .Hepatology, 2019, 70: 563-576. Zhang Yi-Fan, HO Mitchell,Humanization of high-affinity antibodies targeting glypican - 3 in hepatocellular carcinoma. [J] .Sci Rep, 2016, 6: 33878.UsageResearch use onlyConjugatePE-conjugated All DIMA PisclaimerDIMA DisclaimerPE-conjugated activity gluba Biotech. They are all under pater tapplication. Any protein sequencing or reverse engineering attempt is prohibited. We are activity glub generated by DIMA Biotech. They are all under pater tapplication. Any protein sequencing or reverse engineering attempt is prohibited. We are activity scrutinizing all pater tapplication to	Clonality	Monoclonal
Recommended DilutionsFlow Cyt 1:100PurificationFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via glycosyl phosphatid/linositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptid/l peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome. Alternative splicing results in multiple transcript variants. Iprovided by RefSeq, Sep 2009] References] Fu Ying, Urban Daniel J.Mani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[]] .Hepatocellular carcinoma.[]] .Sci Rep, 2016, 6: 33878.UsageResearch use only GonjugateConjugatePE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under protein sequencing or reverse engineering attempt is prohibited. We are activity scrutining all patent application to	Reactivity	Human
DilutionsFlow Cyl 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009] References] Fu Ying, Urban Daniel J.Nain Roger R et al. (Bylcian-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carinoma.[]]. Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell.Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carinoma.[]]. Sci Rep, 2016, 6: 33878.UsageResearch use onlyConjugatePE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA BisclaimerDIMA DisclaimerPE-conjugated actively scrutinizing all patent application to	Applications	Flow Cyt
Formulation & ReconstitutionLiquid]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq. Sep 2009] References] Fu Ying, Urban Daniel J, Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[J] .Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.[J] .Sci Rep, 2016, 6: 33878.UsageResearch use onlyConjugatePE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to actively scrutinizing all patent application to actively scrutinizing all patent application to		Flow Cyt 1:100
ReconstitutionLiquid[]#BS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsCell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via glyccosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq. Sep 2009] References] Fu Ying,Urban Daniel J.Nani Roger R et al. Glypican-3 Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[J] .Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.[J] .Sci Rep, 2016, 6: 33878.UsageResearch use onlyConjugatePE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to bi	Purification	Purified from cell culture supernatant by affinity chromatography
 Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substitute with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi-Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009] References[] Fu Ying, Urban Daniel J.Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[]] Hepatocellular carcinoma.[]] Hepatocellular carcinoma.[]] .Sci Rep, 2016, 6: 33878. Usage Research use only Conjugate PE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibide. We are actively scrutinizing all patent application to 		Liquid PBS with 0.05% Proclin300, 1% BSA
 background Background	Storage & Shipping	Store at 2°C-8°C for 6 months
UsageResearch use onlyConjugatePE-conjugatedAll DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to	Background	composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi- Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009] References[] Fu Ying,Urban Daniel J,Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[J] .Hepatology, 2019, 70: 563-576. Zhang Yi-Fan,Ho Mitchell,Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular
Conjugate PE-conjugated All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to	Usage	
DIMA Disclaimer All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to	-	-
		All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to

Email: info@dimabio.com Website: www.dimabio.com

