

PRODUCT INFORMATION

TargetIL13SynonymsIL-13;NC30Host SpeciesRabbitDescriptionChimeric mAbDeliveryUnder DevelopmentUniprot IDP35225IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytPurificationsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & RecomstitutionStore at 2°C-8°C for 6 monthsStorage & ShippingStore at 2°C-8°C for 6 monthsBackgroundLiquid[DPBS with 0.05% Proclin300, 1% BSABackgroundUnibits the production of pro-inflammatory cytokine is involved in several stages of B-cell matures and activity, thereby inhibits the production of pro-inflammatory cytokine is involved in several stages of B-cell matures and chromokines. This cytokine induced primarity by activated Th2 cells. This cytokine is found to be critical to the pathogenesis of allergen- induced astima but operates through mechanisms independent of IgE and eosinophils. This gene encloses to IL4. [provided by RefSeq. jul 2008]BackgroundResearch use only	Clone ID	1A1
Host SpeciesRabbitDescriptionPE-conjugated Anti-IL13 antibody(1A1), IgG1 Chimeric mAbDeliveryUnder DevelopmentUniprot IDP35225IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionStore at 2°C-8°C for 6 monthsStorage & ShippingStore at 2°C-8°C for 6 monthsBackgroundStore at 2°C-8°C for 6 monthsBackgroundHum Cytabit and GFB and Cytabit and provide a strange activity, thereby inhibits the production of pro-inflammatory cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene elus. This gene enclusets through amechanisms independent of IgE and eosinophils. This gene elus. Lis. Li4, and CSF2 form a cytokine induced asthma but operates through amechanisms independent of IgE and eosinophils. This gene elus. Lis. Li4, and CSF2 form a cytokine pare cluster on chromosome 5q, with this gene pare cluster on chromosome 5q, with	Target	IL13
DescriptionPE-conjugated Anti-IL13 antibody(1A1), IgG1 Chimeric mAbDeliveryUnder DevelopmentUniprot IDP35225IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow CytPurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionStore at 2°C-8°C for 6 monthsStorage & ShippingStore at 2°C-8°C for 6 monthsBackgroundThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes class and chemokines. This cytokine is cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operaes through mechanisms independent of IgE and eosinophils. This gene gene cluster on chromosome 5q, with this gene gene cluster on chromosome 5q, with this gene gene gene cluster on chromosome 5q, with this gene gene cluster on chromosome 5q, with this gene <th>Synonyms</th> <th>IL-13;NC30</th>	Synonyms	IL-13;NC30
DescriptionChimeric mAbDeliveryUnder DevelopmentUniprot IDP35225IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months 	Host Species	Rabbit
Uniprot IDP35225IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionStore at 2°C-8°C for 6 monthsStorage & ShippingStore at 2°C-8°C for 6 monthsBackgroundThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes Igf isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokine is independent of Igf and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene penduction of L4. [provided by RefSeq, Jul 2008]	Description	
IgG typeRabbit/Human Fc chimeric IgG1ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- induced astma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene gene cluster on chromosome 5q, with this gene gene cluster on chromosome 5q, with this ge	Delivery	Under Development
ClonalityMonoclonalReactivityHumanApplicationsFlow CytRecommendedFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene paticularly close to IL4. [provided by RefSeq, Jul 2008]	Uniprot ID	P35225
ReactivityHumanApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	lgG type	Rabbit/Human Fc chimeric IgG1
ApplicationsFlow CytRecommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	Clonality	Monoclonal
Recommended DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid_PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine is found to be critical to the pathogenesis of allergen- inhibits the production of pro-inflammatory cytokine is of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	Reactivity	Human
DilutionsFlow Cyt 1:100PurificationPurified from cell culture supernatant by affinity chromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 months This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	Applications	Flow Cyt
FormulationchromatographyFormulation & ReconstitutionLiquid[]PBS with 0.05% Proclin300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IJE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene lL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]		Flow Cyt 1:100
ReconstitutionLiquid PBS with 0.05% Proclim300, 1% BSAStorage & ShippingStore at 2°C-8°C for 6 monthsThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	Purification	
BackgroundThis gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]		Liquid PBS with 0.05% Proclin300, 1% BSA
Backgroundproduced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]	Storage & Shipping	Store at 2°C-8°C for 6 months
· · · · · · · · · · · · · · · · · · ·	Background	produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen- induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul
osuge Acocaren ase only	Usage	Research use only
Conjugate PE-conjugated	Conjugate	PE-conjugated

