

PRODUCT INFORMATION

Clone ID DM148 IL2RA **Target**

IL2RA;CD25;p55;IL2-RA;IL-2-RA **Synonyms**

Host Species

PE-conjugated Anti-IL2RA antibody(DM148); Rabbit mAb Description

Delivery 3-4 weeks **Uniprot ID** P01589 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Liquid PBS with 0.05% Proclin300, 1% BSA

Storage & Shipping Store at 2°C-8°C for 6 months

> The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains; together with the common gamma chain (IL2RG); constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor; while homodimeric lasts (IL2RB) chains produce a modium officials. beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein; soluble IL2RA has been isolated and determined


to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated; but the significance of each is presently unknown. Mutations in this gene are associated

> Email: info@dimabio.com Website: www.dimabio.com

with interleukin 2 receptor alpha deficiency.

Usage Research use only

Conjugate PE-conjugated



/+86-400-006-0995(China)

