

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM71
<b>Target</b>	Mesothelin
<b>Synonyms</b>	MSLN; Mesothelin; MPF
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-mesothelin antibody(DM71), Rabbit mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	Q13421
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	This gene encodes a preproprotein that is proteolytically processed to generate two protein products; megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiating factor functions as a cytokine that can stimulate colony formation of bone marrow megakaryocytes. Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas; ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants; at least one of which encodes an isoform that is proteolytically processed.
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

