

PRODUCT INFORMATION

Clone ID **DMC493** MUC1 **Target**

ADMCKD; ADMCKD1; CA 15-3; CD227; EMA; H23AG; KL-6; MAM6; MCD; MCKD; MCKD1; MUC-1; MUC-1/SEC; MUC-1/X; MUC1/ZD; PEM; PEMT; PUM **Synonyms**

Host Species

PE-conjugated Anti-MUC1 antibody(DMC493), **Description**

IgG1 Chimeric mAb

3-4 weeks Delivery P15941 **Uniprot ID**

Rabbit/Human Fc chimeric IgG1 IgG type

Monoclonal Clonality Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Liquid

□PBS with 0.05% Proclin300, 1% BSA Reconstitution

Store at 2°C-8°C for 6 months Storage & Shipping

> This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role

in forming protective mucous barriers on

epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the discount of the surfaces of many different tissues included in the surfaces of many different tissues including lung; breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric

complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression; aberrant intracellular localization; and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results

> Email: info@dimabio.com Website: www.dimabio.com

in multiple transcript variants.

Usage Research use only Conjugate PE-conjugated

