

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DMC222 |
| Target | PCSK9 |
| Synonyms | FH3; FHCL3; HCHOLA3; LDLCQ1; NARC-1; NARC1; PC9 |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-PCSK9 antibody(DMC222); IgG1 Chimeric mAb |
| Delivery | 3-4 weeks |
| Uniprot ID | Q8NBP7 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| Storage & Shipping | Store at 2°C-8°C for 6 months |
| Background | <p>This gene encodes a member of the subtilisin-like proprotein convertase family; which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver; intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants.</p> |
| Usage | Research use only |
| Conjugate | PE-conjugated |

