

## PRODUCT INFORMATION

<b>Clone ID</b>	DMC389
<b>Target</b>	VWF
<b>Synonyms</b>	F8VWF; VWD
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-VWF antibody(DMC389); IgG1 Chimeric mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	P04275
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	This gene encodes a glycoprotein involved in hemostasis. The encoded preproprotein is proteolytically processed following assembly into large multimeric complexes. These complexes function in the adhesion of platelets to sites of vascular injury and the transport of various proteins in the blood. Mutations in this gene result in von Willebrand disease; an inherited bleeding disorder. An unprocessed pseudogene has been found on chromosome 22.
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

