

PRODUCT INFORMATION

Target	S1 protein NTD
Synonyms	S1 protein NTD;Spike protein S1 NTD;BetaCoV S1-NTD;COVID-19
Description	Recombinant SARS-CoV-2 (2019-nCoV) S1 protein NTD with C-terminal mouse Fc tag
Delivery	In Stock
Uniprot ID	P0DTC2
Expression Host	HEK293
Tag	C-Mouse Fc Tag
Molecular Characterization	S1 protein NTD(Ser13-Leu303) mFc(Pro99-Lys330)
Molecular Weight	The protein has a predicted molecular mass of 59.1 kDa after removal of the signal peptide.The apparent molecular mass of S1-NTD-mFc is approximately 70-100 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.
Usage	Research use only
Conjugate	Unconjugated





Figure 1. SARS-CoV-2 (2019-nCoV) S1 protein NTD, mFc Tag on SDS-PAGE under reducing condition.

SARS-CoV-2 (2019-nCoV) S1 NTD, mFc Tagged protein ELISA




Figure 2. ELISA plate pre-coated by 2 $\mu\text{g/ml}$ (100 $\mu\text{l/well}$) Human AXL, His tagged protein (PME100070) can bind Human NTD, mFc Tagged protein (PME100492) in a linear range of 0.006-0.16 $\mu\text{g/ml}$.

